

Original Article

## Isolation, Identification and Phylogeny Analysis of Fungi from Motichoor Ladoos belonging to the Hadapsar Village of Pune district, Maharashtra, India

<sup>1</sup>Sonali Santosh Kadam, <sup>2</sup>Javalkar Pradnya Santosh, <sup>3</sup>Awale Shreeya Kishor, <sup>4</sup>Mulani Aamna Nasir, <sup>5</sup>Latthe Vaishnavi Aba, <sup>6</sup>Sakshi Madhukar Nanwate, <sup>7</sup>Vaishnavi Tanaji Ingale, <sup>8</sup>Sakshi Sutar, <sup>9</sup>Chaitali Autade, <sup>10</sup>Atharva Vivekanand Sasane, <sup>11</sup>Nikhil Nanasaheb Pawar, <sup>12</sup>Shilpa Lokhande, <sup>13</sup>Smita Karne, <sup>14</sup>Rajesh Dhakane, <sup>15</sup>Sapna Rambahadur Thapa

<sup>1</sup>Department of Botany, R.P. Gogate and R. V. Jogalekar College, Ratnagiri, Maharashtra, India.

<sup>2,3,4,5,6,7,8,9,10,11,13,14</sup>Department of Microbiology, Jayawantrao Sawant Commerce and Science College of, Hadapsar, Pune, Maharashtra, India.

<sup>12</sup>Department of Microbiology, Sant Gadge Baba Amravati University, Amravati, Maharashtra, India.

<sup>15</sup>Department of Microbiology, S.M. Joshi College, Hadapsar, Pune, Maharashtra, India.

### Article Info

#### Article history:

Received: January 1, 2024

Accepted: January 11, 2024

Published: January 14, 2024

Corresponding Author: Rajesh Dhakane

Email: [rajeshdhakane001@gmail.com](mailto:rajeshdhakane001@gmail.com)

**Keywords:** BLAST, GenBank, ITS, *Alternaria burnsii*

### Abstract

The purpose of this study is to identify and isolate fungus from motichoor ladoos from the Hadapsar region in Pune area of Maharashtra, India. Sequence analysis, BLAST searches, PCR amplification, sequencing of the internal transcribe spacer region (ITS) were used to characterize the isolated fungi. After BLAST analysis, the obtained sequences were compared to the sequences from GenBank, the corresponding fungal isolate was identified at the species level (*Alternaria burnsii*).

©Author(s). This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-nc-sa/4.0/) that permits noncommercial use of the work provided that credit must be given to the creator and adaptation must be shared under the same terms.

### 1. Introduction:

Fungi are eukaryotic in nature. They might take the shape of molds, yeast, or a mixture of the two. Certain fungi can induce allergic, systemic, superficial, cutaneous, or subcutaneous illnesses. Yeasts are microscopic fungus made up of single cells that proliferate by budding. The chitinous microfibrils contained in a matrix of tiny polysaccharides, proteins, lipids, inorganic salts, and pigment make up the layered structure that makes up the fungal cell wall. This structure gives the enclosing protoplast shape and skeletal support.

The morphological traits of *Alternaria burnsii* CBS 107.38, such as conidia dimension, color, and septa (longitudinal and transverse); conidiophore type, size, and septa; sporulation patterns (in chains or solitary; branched or unbranched); and cultural traits, have all been taken into consideration when classifying this species [1, 2-5]. The fungus *Alternaria* is responsible for aspergillosis, candidiasis, asthma, and upper respiratory tract infections and their presence in food is needed to be detected to protect health of people across.

In this work, the isolation, identification and phylogeny analysis of fungi from Motichoor ladoos

belonging to the Hadapsar village of Pune district, Maharashtra, India was studied based on its molecular analysis.

## 2. Materials and Method:

### 2.1 Isolation, DNA Extraction, PCR and Sequencing:

Fungi species was isolated from Motichoor ladoos collected from one of the Sweet Homes from Hadapsar, Maharashtra, India on Potato Dextrose Agar [PDA] (hereafter, will be written as PDA) at  $28\pm 2^\circ\text{C}$  for 8 days and maintained at  $4^\circ\text{C}$  for culturing purposes. The morphological characters of fungi were visualized by Lactophenol Blue staining technique [6].

The identification of isolates was carried out at the sequencing facility of National Centre for Microbial Resource (NCMR), National Centre for Cell Science, Pune. At the facility, genomic DNA was isolated by the standard phenol/chloroform extraction method [7], followed by PCR amplification of the ITS regions using universal primers ITS1 [5'-TCC GTA GGT GAA CCT GCG G -3'] and ITS4 [5'-TCC TCC GCT TAT TGA TAT GC -3'].

The amplified ITS PCR product was purified by PEG-NaCl precipitation and directly sequenced on an ABI® 3730XL automated DNA sequencer (Applied Biosystems, Inc., Foster City, CA) as per manufacturer's instructions. Essentially, sequencing was carried out from both ends so that each position was read at least twice. Assembly was carried out using Lasergene package followed by NCBI BLAST against sequences from type material for tentative identification [8].

### 2.2. Phylogenetic Analysis:

They phylogenetic analysis of the obtained fungal sequence was performed using Molecular Evolutionary Genetic Analysis X (MEGA X) [9] with Neighbour-Joining (NJ) method of analysis and a bootstrap support of 1000 replicates by taking the sequences from NCBI database showing highest similarities.

## 3. Result

The sample of fungi was processed for identification on the basis of ITS region (table 1). The obtained sequence was then submitted to NCBI with accession number QQ928954.1. The identification was performed using NCBI database. The reliability in identification is restricted to not only availability but also the homology extent revealed by approximately 550 base pairs (bp)

sequence of the sample under study with its closest neighbor in the database

The comparison of the sequence with NCBI database (NCBI-BLASTn (<http://blast.ncbi.nlm.nih.gov/>)) [10] showed that the fungi under study is *Alternaria burnsii* with 99.61% similarity with other sequences in the database. The fungal sample under study (A JAN 23 096) showed cluster (in a single clade) with NCBI sequences KP 124420.1, *Alternaria burnsii* strain CBS 107.38, and KP136019.1 *Alternaria burnsii* strain CBS 107.38. As well, threads of fungi reported in Lactophenol Blue staining [figure 1].

## 4. Discussion:

The isolate from Motichoor ladoos collected from Hadapsar, Maharashtra, India was similar with those described as *Alternaria burnsii*. This finding was supported by Paul et al. [11] who reported that the fungal isolates from Bangladesh Agricultural Research Institute (BARI), Bangladesh pumpkin seeds were similar to *Alternaria bursnii*. The ITS sequences of fungal isolates from Bangladesh showed over 100% sequence similarities with *A. burnsii* [11]. Similarly, the fungus specimen with Genbank accession no. QQ928954.1 used in our study reported 99.61% similarity with the NCBI sequence database. The difference of 0.39% sequence similarity was reported in both studies with the NCBI sequence database.

## 4. Conclusion:

The fungal species present in the Motichoor ladoos can be identified by using Internal Transcribed Spacer 1 [ITS up to species level using NCBI sequence database. However, the species identification is not robust and reliable since there is variation in the sequence similarities from different geographical locations (India and Bangladesh).

### Acknowledgement:

The authors are thankful to Dr. Sandip Gawate, I/C Principal, Jayawantrao Sawant Commerce and Science College, Hadapsar, Pune, Maharashtra, India for providing laboratory facilities to conduct the research work.

**Author contribution:** SSK, JPS, ASK, MAN, LVA, SMN, VTI: Developed and idea and conducted experimental work, SS, CA, AVS, NNP, SL, SK, ST: Wrote the manuscript, RD: Verified the content.

**Competing interest:** Authors declare that no competing interest exists.

**Ethical statement:** Since the work does not violate ethical laws, no ethical permission required.


**Grant Support Details:** The authors did not receive funding for this work from any agency.

**References:**

1. Simmons EG. *Alternaria: an identification manual*. Utrecht: CBS Fungal Biodiversity Centre; 2007.
2. Sung GH, Sung JM, Hywel-Jones NL, Spatafora JW. A multi-gene phylogeny of Clavicipitaceae (Ascomycota, Fungi): identification of localized incongruence using a combinational bootstrap approach. *Mol Phylogenet Evol.* 2007;44(3):1204-23. doi: [10.1016/j.ympev.2007.03.011](https://doi.org/10.1016/j.ympev.2007.03.011), PMID [17555990](https://pubmed.ncbi.nlm.nih.gov/17555990/).
3. Liu YJ, Whelen S, Hall BD. Phylogenetic relationships among ascomycetes: evidence from an RNA Polymerase II Subunit. *Mol Biol Evol.* 1999;16(12):1799-808. doi: [10.1093/oxfordjournals.molbev.a026092](https://doi.org/10.1093/oxfordjournals.molbev.a026092), PMID [10605121](https://pubmed.ncbi.nlm.nih.gov/10605121/).
4. Hong SG, Cramer RA, Lawrence CB, Pryor BM. Alt a 1 allergen homologs from *Alternaria* and related taxa: analysis of phylogenetic content and secondary structure. *Fungal Genet Biol.* 2005;42(2):119-29. doi: [10.1016/j.fgb.2004.10.009](https://doi.org/10.1016/j.fgb.2004.10.009), PMID [15670710](https://pubmed.ncbi.nlm.nih.gov/15670710/).
5. Berbee ML, Pirseyedi M, Hubbard S. Cochliobolus phylogenetics and the origin of known, highly virulent pathogens, inferred from ITS and glyceraldehyde-3-phosphate dehydrogenase gene sequences. *Mycologia.* 1999;91(6):964-77. doi: [10.1080/00275514.1999.12061106](https://doi.org/10.1080/00275514.1999.12061106).
6. Sambrook J, Fritsch EF, Maniatis T. *Molecular Cloning a laboratory manual*. New York: Cold Spring Harbor Laboratory Press; 1989.
7. Woudenberg JH, Groenewald JZ, Binder M, Crous PW. *Alternaria* redefined. *Stud Mycol.* 2013;75(1):171-212. doi: [10.3114/sim0015](https://doi.org/10.3114/sim0015), PMID [24014900](https://pubmed.ncbi.nlm.nih.gov/24014900/).
8. Paul NC, Kim WK, Woo SK, Park MS, Yu SH. Diversity of endophytic fungi associated with *Taraxacum coreanum* and their antifungal activity. *Mycobiology.* 2006;34(4):185-90. doi: [10.4489/MYCO.2006.34.4.185](https://doi.org/10.4489/MYCO.2006.34.4.185), PMID [24039496](https://pubmed.ncbi.nlm.nih.gov/24039496/).
9. Woudenberg JH, Groenewald JZ, Binder M, Crous PW. *Alternaria* redefined. *Stud Mycol.* 2013;75(1):171-212. doi: [10.3114/sim0015](https://doi.org/10.3114/sim0015), PMID [24014900](https://pubmed.ncbi.nlm.nih.gov/24014900/).
10. Reuveni R, Agapov V, Reuveni M, Raviv M. Effects of foliar sprays of phosphates on powdery mildew (*Sphaerotheca pannosa*) of roses. *J Phytopathol.* 1994;142:331-7.
11. Wang XX, Chen J, Wang B, Liu LJ, Huang X, Ye ST et al. First report of anthracnose on *Boehmeria nivea* caused by *Colletotrichum higginsianum* in China. *Plant Dis.* 2011;95(10):1318-. doi: [10.1094/PDIS-04-11-0356](https://doi.org/10.1094/PDIS-04-11-0356), PMID [30731673](https://pubmed.ncbi.nlm.nih.gov/30731673/).

**Cite this article as**

Kadam SS, Javalkar PS, Awale SK, Nasir MA, Latthe VA, Nanwate SM, Ingale VT, Sutar S, Autade C, Sasane AV, Pawar NN, Lokhande S, Karne S, Dhakane R, Thapa SR. Isolation, identification and phylogeny analysis of fungi from Motichoor ladoos belonging to the Hadapsar village of Pune district, Maharashtra, India. *Int. J. Micro. Sci.* 2024; 5(1), 1-5.



**Figure 1: The visualized fungi under 45X objective during Lactophenol Blue Staining**

**Table 1: Isolated Fungi gene sequence**

---

**Amplicon sequence**

---

>A\_JAN\_23\_096 *Alternaria burnsii*

CTTCCGTAGGGTGAACCTGCGGAGGGGATCATTACACAAATATGAAGGCGGGCTGGAATCTCTCGGGGTTACAGCCTTGCTGAATTATTCACCCTTGCTTTTGGCGTA  
 CTTCTTGTTTCCTTGGTGGGTTGCGCCACCACTAGGACAAACATAAACCTTTTGAATTGCAATCAGCGTCAGTAACAAATTAATAATTACAACCTTTCAACAACGGATC  
 TCTTGGTTCTGGCATCGATGAAGAACGCAGCGAAATGCGATAAGTAGTGTGAATTGCAGAATTCAGTGAATCATCGAATCTTTGAACGCACATTGCGCCCTTTGGTA  
 TTCCAAAGGGCATGCCTGTTGAGCGTCATTTGTACCCTCAAGCTTTGCTTGGTGTGGGCGTCTTGTCTAGCTTTGCTGGAGACTCGCCTTAAAGTAATTGGCAG  
 CCGGCCTACTGTTTTGGAGCGCAGCACAAAGTCGCACTCTCTATCAGCAAAGGTCTAGCATCCATTAAGCCTTTTTTCAACTTTTGACCTCGGGATCAGGTAGGGAT  
 ACCCGCTGAACCTAAGCATATCAATAGGCGGAGGAA

---




Figure 2: Phylogenetic tree constructed using MEGA 10 [35].